

T1.1D - Psychometric validation of the Computer Proficiency Questionnaire (CPQ-Brazil) adapted to Brazilian Portuguese

Gabrieli Pereira da Cruz¹

gabrielicruz@usp.br

Paula Costa Castro¹

castro@ufscar.br

Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador São Carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

Abstract.

The rapid demographic transition and technological advancement emphasize the need for older adults to adapt to the digital world. However, validated tools to assess their technological skills and support digital inclusion are limited. This study aims to validate the Brazilian version of the Computer Proficiency Questionnaire, adapted to assess technological literacy in older adults. A methodological study is being conducted with 350 participants aged 60 and above, recruited nationwide. Participants completed a socio-demographic questionnaire, and the validation process includes assessments of reproducibility, reliability, and both exploratory and confirmatory factor analyses. To date, 226 older adults (mean age= 71.19 years, SD= 6.36) have participated, with 71.2% identifying as female. The average years of schooling was 12.80 (SD = 6.07), with a median of 12 years and a range of 0 to 35 years. Most participants were from the South (77.73%), followed by the Southeast (19.91%). Responses were collected via paper (42%) and online (58%). Regarding internet use, 59.3% accessed it more than once a day, 3.5% had no access, and 6.6% had access but did not use it. Most participants (68.1%) completed the questionnaire unaided. Correlation analyses revealed a weak negative relationship between age and technological proficiency ($r = -0.23$) and a moderate positive correlation with years of schooling ($r = 0.62$). The highest proficiency score was in 'Computer Basics' (mean= 3.60), while 'Diary/Calendar' had the lowest (mean= 2.33). Final results will clarify the tool's applicability.

Keywords. Older people; Education and training; Technology; Questionnaire; Psychometric validation.

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T1.2D - Lung opacities Multimodal prognostic evaluation using Artificial Intelligence

Johanna Elisabeth Rogalsky¹
jerogalsky@usp.br

Lucas Ferrari de Oliveira³
lferrari@inf.ufpr.br

Marcel Koenigkam Santos²
marcelk46@fmrp.usp.br

Paulo Mazzoncini de Azevedo Marques¹
pmarques@fmrp.usp.br

¹Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
São-carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

²Faculdade de Medicina de Bauru - Universidade de São Paulo, Alameda Dr. Octávio Pinheiro
Brisolla, 9-75 - Vila Regina, 17012-230, Bauru - SP – Brazil

³Departamento de Informática (DInf), Universidade Federal do Paraná - UFPR, R. Evaristo F.
Ferreira da Costa, 383-391 - Jardim das Américas, 81530-090, Curitiba - PR – Brasil

Abstract. Interstitial lung diseases (ILDs) compose a group of more than 200 distinct pathological processes. Despite this, they may share symptoms and features on examinations, making their differential diagnosis more difficult. The diagnosis uses the computed tomography (CT) examination, which is also present in patient management, being part of the prognosis and evaluation of the disease and the treatment efficacy. However, the visual analysis of the exams performed by radiologists is a complex task, which can be tiring and result in intra- and inter-observer variation regarding radiological and diagnostic findings. In this context, computational support has become a part of this evaluation. By associating this quantification and clinical data with a clinical outcome, radiomics has opened new avenues for subtyping and phenotyping these lung diseases. This way, computational tools able to infer the clinical outcome can assist in the diagnosis, but mainly in the longitudinal management of the patient, given an association of quantification of features extracted from CT images and data from additional exams. Retrospectively combining clinical exams and data from a group of patients with ILD enables building a model that infers the clinical outcome. Based on the above, this project aims to investigate computational models that use image registration, machine learning, and deep learning techniques, so that they can be used to perform radiomic association of multimodal data and enable the diagnostic and prognostic evaluation of patients with ILD.

Keywords. Radiomics. Interstitial Lung Diseases. Machine Learning. Deep Learning.

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T1.3D - Assessment of the Radiomics Pipelines for Histological Classification of Non-Small Cell Lung Cancer from Molecular Images of [18F]FDG-PET/CT

Alexandre-Santos, L.^{2,3}

leonardosantos.mp@usp.br

Wichert-Ana, L.^{1,2,3}

lwichert@fmrp.usp.br

1Ribeirão Preto Medical School - University of São Paulo, Ribeirão Preto, BRAZIL, 2Nuclear Medicine and PET/CT Section, The Clinical Hospital, Ribeirão Preto, BRAZIL, 3The Interunits Bioengineering Posgraduation Program, São Carlos School of Engineering, University of São Paulo, São Carlos SP, BRAZIL.

In the staging of non-small cell lung cancer (NSCLC) cases, histological assessment is gold standard in characterization of type tumor. However, viable tissue samples for evaluation is challenging, what can be more explored with radiomic techniques to aid in this classification by highlighting areas of histological heterogeneity. This study aimed to evaluate the performance of various machine learning pipelines for classifying NSCLC cases using [18F]FDG-PET-CT images. **Materials and methods:** We investigated 135 PET images of patients diagnosed with NSCLC of type adenocarcinoma (ADS) and squamous cell carcinoma (SCC), confirmed through histopathological assessment. The images were manually segmented by a nuclear medicine physician and extracted 2.082 radiomic features that could potentially contribute to histological classification. We used three feature selection methods (F-Score (ANOVA), Mutual Information (MI), and Principal Component Analysis (PCA)) along with six classification methods (Random Forest, Support Vector Machine, k-Nearest Neighbors, Gaussian Naïve Bayes, Decision Tree, and XGBoost) to build machine learning pipelines capable of differentiating between the two types of NSCLC. The model complexity and area under the curve (AUC) were used as metrics to assess the performance of the methods. **Results:** The eighteen pipelines of classification assessed evidenced good performance for XGBoost+ANOVA and MI (AUC > 0.9 to test) and increased performance for the pipeline with PCA for dimensional reduction. **Conclusion:** The radiomic machine learning methods for classifying NSCLC cases have demonstrated good performance in distinguishing [18F]FDG-PET/CT images of ADS and SCC when implemented with feature selection methods ANOVA and MI, however, compromised for the PCA method.

Keywords. Nuclear Medicine, NSCLC, PET-CT, Radiomics, Machine learning methods.

T2.1M - Automatic detection and segmentation of tumors in chest computed tomography exams to aid in the diagnosis and radiomics of lung cancer

Douglas Samuel Gonçalves¹

douglas.samuel@usp.br

Paulo Mazzoncini de Azevedo Marques¹

pmarques@fmrp.usp.br

*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
Sãocarlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

Abstract. Lung cancer is the type of cancer with one of the highest mortality rates in Brazil, and the highest rate in the world. With the advance of technology, systems have been created that work extensively in the medical field to help analyze these tumors in patient examinations, known as computer-aided diagnosis (CAD). The process of radiomics is an extension of CAD systems, which includes transforming images from medical examinations into data that can be interpreted by algorithms, viewing images as numbers and sequences of intensities that mathematically determine what these lesions are in the examinations, allowing important characteristics to be extracted for classifying the lesion. Convolutional neural networks (CNN) are deep learning methods used primarily in image classification, where the image is processed by multiple layers that apply filters and generic functions to extract characteristics from these images and classify them. Thus, the aim of this research is to use algorithms and methods for segmenting lung lesions and structures pertinent to the lung to create a database with images of tumors and structures that can generate a false positive in the clinical evaluation of the exam and use this database to train a CNN type neural network and assess the feasibility of creating an automated segmentation algorithm for lung lesions.

Keywords. Convolutional Neural Networks; Image Segmentation; Deep Learning; Radiomics; Automatic Segmentation.

T2.2M - Study of machine learning models for texture characterization of vertebral bodies in magnetic resonance imaging and correlation with compression fractures.

Laura Teresa Ricoboni¹

lauraricoboni@usp.br

Paulo Mazzoncini de Azevedo Marques¹

pmarques@fmrp.usp.br

Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador São-carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

Abstract.

Osteoporosis is a chronic osteometabolic disease characterized by reduced bone mineral density (BMD), which leads to an increased risk of fractures. The latest definitions of osteoporosis introduce the concept of bone quality, encompassing a set of structural and compositional characteristics of bone responsible for maintaining skeletal resistance to fractures. According to the World Health Organization (WHO), fragility fractures are those that occur as a result of low-energy trauma, resulting from a reduction in bone strength and reflecting skeletal fragility under biomechanical stress. In addition to causing loss of mobility, institutionalization, and increased risk of mortality, these fractures increase healthcare costs due to their high prevalence in individuals with multiple comorbidities. In the spine, compression fractures occur when the load on the vertebral body exceeds its stability, resulting in bone collapse. Studies indicate that people with osteoporotic vertebral fractures have a significantly higher risk of subsequent fractures. The measurement of bone mass through dual-energy X-ray absorptiometry (DXA) is considered the “gold standard” for the diagnosis of osteoporosis. Although validated in clinical trials and widely used for diagnosis and monitoring of the disease, DXA has limitations in predicting fracture risk in osteoporotic patients. Therefore, this project aims to develop machine learning models that complement the diagnosis performed by DXA by investigating biomarkers and characterizing the texture of non-fractured vertebral bodies, correlating them with vertebral bodies that present compression fractures.

Keywords. Osteoporosis, Bone fractures, Dual-energy X-ray absorptiometry (DXA), Machine learning models.

Acknowledgments.

We would like to thank the Faculty of Medicine of Ribeirão Preto, the School of Engineering of São Carlos and the Institute of Chemistry of São Carlos of the University of São Paulo, the Center for Imaging Sciences and Medical Physics of the Hospital das Clínicas of Ribeirão Preto, as well as the funding agency CAPES for the financial support.

T2.3D - Development of a protocol for the evaluation of the effects of non-immersive virtual reality practice on the motor control of young adults

Bruna Carolina Mania Duarte¹

brunaduarte@usp.br

Alexandre Fonseca Brandão²

alexandre.brandao@puc-campinas.edu.br

Daniela Godoi-Jacomassi³

danielagodoij@ufscar.br

¹*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador São-carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

²*Research Professor at the Polytechnic School of the Pontifical Catholic University of Campinas, R. Professor Dr. Euryclides de Jesus Zerbini, 1516 - Parque Rural Fazenda Santa Cândida, 13087-571, Campinas – SP, Brazil.*

³*Dinâmica – Motor Behavior Laboratory, Department of Physical Education, Federal University of São Carlos, Rod. Washington Luis Km235, 13565-905, São Carlos – SP, Brazil.*

Abstract.

Virtual Reality (VR) can potentially contribute to health as an alternative to traditional interventions. Few studies examined young adults and the effects of these practices on motor control (MC). Moreover, there aren't studies that compared the adaptations resulting from the practice of VR (non-immersive) in this population that is considered a standard for comparison. Therefore, the primary objective was the development of a practice and assessment protocol that allows the investigation of the effects of VR practice and real practice (RP) on MC in young adults. The secondary objective was the application of the protocol developed to evaluate its applicability and feasibility. For this, (a) a protocol of practice with activities to be carried out in VR and RP was developed, as well as an evaluation protocol to compare the effects of practice with VR and RP on the MC of young adults; and (b) an application study was carried out, in which two young adults participated. The results showed (a) the process of developing the practice protocol, which included the practice of games in both practices during 18 sessions of 20 minutes, three times a week, and the assessment protocol that included assessments of postural control, heart rate variability and cortical activation before and after the intervention and one week after the intervention; and that (b) the protocol developed can be applied and proved to be viable to compare the effects of practices with VR and RP on the MC of young adults.

Keywords. Virtual reality; Motor control; Young adults.

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T2.4M - Knowledge, perception and attitude regarding artificial intelligence among dentists: A systematic review

Helena Cristina Aguiar¹

hcaguiar@usp.br

Kaíssa da Cunha Lima¹

kaissalima@usp.br

Camila Tirapelli¹

catirapelli@forp.usp.br

Brigitte Grosgeat²

brigitte.grosgeat@univ-lyon1.fr

¹*Graduate Program in Oral Rehabilitation, Department of Dental Materials and Prosthesis, Ribeirão Preto School of Dentistry, University of São Paulo, Café Avenue S/N, Ribeirão Preto 14040-904 – SP - Brazil;*

²*Faculté d'Odontologie, Laboratoire des Multimatériaux et Interfaces, UMR CNRS 5615, Pôle d'Odontologie, Hospices Civils de Lyon - Lyon - France;*

Abstract. Applications of artificial intelligence in dentistry are advancing rapidly, enhancing treatment precision, reducing consultation time, and aiding in the diagnosis and prognosis of dental conditions. However, as these technologies are relatively new, several challenges arise in clinical implementation, including ethical dilemmas and the need for professional training. This review aims to understand dentists' knowledge, attitude and perception of artificial intelligence. An electronic search was conducted across PubMed, Embase, Scopus, Web of Science, Lilacs, ACM Digital Library, and Google Scholar databases, resulting in 3,132 articles, of which 630 duplicates were excluded. The remaining 2,502 articles underwent a double-blind screening process, and 48 articles were selected for full-text review. A total of 23 articles were included from the database search, and 6 were included through citation searching. The articles were assessed for risk of bias using the Joanna Briggs Institute (JBI) Checklist for Analytical Cross-Sectional Studies and were rated according to the GRADE scale for quality of evidence. Studies have shown a positive association between dentists' perceptions and the application of artificial intelligence (AI) in dentistry. Despite significant variation in professionals' knowledge levels regarding AI, most studies highlight the need to invest in training for both students and practicing professionals. Participants remain concerned with ethical issues such as data privacy, liability for errors, and algorithm transparency. Therefore, while optimism toward AI has been noted, advancements in education and ethical frameworks are essential for the successful implementation of this technology in clinical practice.

Keywords. Artificial intelligence; Dentistry; Attitude; Perception; Awareness; Systematic review

T3.1D - Effect of pH and Ionic Strength on the Antimicrobial Activity of Rhamnolipids Against *Staphylococcus aureus*

Amanda Mezete Gouveia¹

amgouveia@usp.br

Prof^a. Dr^a. Marcia Nitschke¹

nitschke@iqsc.usp.br

Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
São-carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

Abstract.

Waterborne and foodborne diseases (WFD) are among the leading causes of mortality in developing countries, with bacteria being the primary pathogens involved. Among these, *Staphylococcus aureus* is notable for causing diseases through infection and food poisoning, resulting from consuming contaminated food. The increasing bacterial resistance to conventional antibiotics, combined with the interest in more sustainable antimicrobials, has driven the search for alternative control methods. In this context, the use of biosurfactants (BS), such as rhamnolipids (RL), emerges as a promising alternative due to their low toxicity, biodegradability, and production from renewable sources. This study aimed to evaluate the effect of pH and ionic strength on the antimicrobial activity of RL against planktonic and sessile forms of *S. aureus*. The results indicated that increasing the pH from 5 to 8 raised the minimum inhibitory concentration (MIC) from 78.12 mg/L to over 2500 mg/L. Moreover, the presence of NaCl enhanced the antimicrobial activity of RL at all tested pH values, reducing MIC values. Planktonic cells of *S. aureus* were more sensitive to RL treatment than biofilms. Despite the greater resistance of biofilms, the presence of RL was able to reduce the number of viable cells by up to 3.8 logarithmic units. It is concluded that both ionic strength and pH significantly influence the antimicrobial activity of RL against *S. aureus*, suggesting its potential application in acidic and salty foods.

Keywords: Ionic strength. Biosurfactant. Biofilms. Rhamnolipid. *S. aureus*.

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T3.2D - Printability of bioinks: a consolidated definition for future prediction models

Matheus A A Resende¹

matheus.a.resende@usp.br

Andres Vercik¹

avercik@usp.br

*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
Sãocarlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

Abstract. Printability is a term widely used nowadays for research on bioinks for additive manufacturing. Despite of this, it seems to be an absence of a common and universally accepted interpretation of the term, which can be source of misunderstanding among researchers, whereas some of them use printability and shape fidelity or shape accuracy indistinctly, other authors take into account different aspects of the printing process, such as, shape fidelity, extrudability, shape accuracy, including them as factors that, combined, lead to a more complete concept. In any case, and based on works reported in literature, a consensus is still lacking. In the case of bioprinting, cell viability is an extremely important factor to be included. This impacts any study on printability, since a partial view it would be based on a partial view of this property. For instance, studies relating rheological properties and printability and the prediction of this property on the physical properties of the inks will suffer of a generality as poor as the definition of the target property, independently of the methodology (physical modeling, machine learning, etc.). Thus, this scenario avoids any prediction of the complete manufacturing process. In this work a literature review between the years 2011 and 2023 on the concept of printability, following the evolution of the term and how it has been increasingly applied in the area. By using data science and machine learning tools, particularly text mining and text analytics, we can identify some cluster ideas leading to a general definition with greater acceptance in a wide range of different areas of application of additive manufacturing. Based on this we propose more general definition of printability in order to build such consensus.

Keywords. Printability; Bioinks; Bioprinting; Cell Viability; Predictive Models

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001

T3.3M - Graphene oxide reduction process: monitoring and modeling of reaction kinetics

Alessandra Maria Garcia Barbosa¹

alessandra.g@usp.br

Andrés Vercik¹

avercik@usp.br

Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador São carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

Abstract.

Graphene is of extreme importance due to its unique physical properties such as mechanical strength, thermal and electrical conductivity and transparency, among others, making this material ubiquitous in science. Unfortunately, obtaining this material is complex and, because of that, considerable efforts are devoted in finding more efficient methods for large-scale and low-cost production of graphene. A viable alternative is the chemical reduction of graphene oxide. Depending on the specific application, a material with properties similar to those of the graphene can be obtained. For instance, by the reduction process the electrical conductivity of the graphene oxide, which is an insulator, can be increased resulting in conductive reduced graphene oxide. This changing in the electrical conductivity can be also used for monitoring the kinetics of the reduction process. Despite the importance of the chemical reduction of graphene oxide, there is a lack of studies of the reaction kinetics of this process, whose description would allow its optimization both in time and in cost. This work aims to show a novel scheme for real time and on-line electrical monitoring of the graphene oxide chemical reduction process. The method is tested in different reduction conditions. The electrical conductivity of the reduction solution is measured continuously during the process. A mathematical model is also proposed to determine the reduced graphene oxide concentration from conductivity according to the following expression: $m = \exp((\kappa - c)/a) - b$, where a , b and c are fitting parameters. This model was validated with other expressions found in literature for the conductivity of electrolyte solutions.

Keywords. Reduction; Kinetics; Mathematical model.

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T4.1D - Nonlinear Analysis by Phase Space Reconstruction in Voice Signals with Benign Laryngeal Lesions.

Mariana Ferreira Gonçalves¹

mariana.ferreira.goncalves@usp.br

Lídia Cristina da Silva Teles^{1,2}

lidiactm@usp.br

¹*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador São Carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

²*Dept. of Speech, Language and Hearing Disorders, Faculty of Dentistry of Bauru, University of São Paulo, Brazil.*

Objective: Describe Phase Space Reconstruction (PSR) in voice signals with benign laryngeal lesions.

Method: 106 voice signals were analyzed from patients at FOB-USP, including 43 with nodules (41 women, 2 men, ages 18–50, average 32 ± 8 years), 40 with cysts (39 women, 1 man, ages 18–47, average 31 ± 8 years), and 23 with sulcus (13 women, 10 men, ages 22–48, mean 34 ± 9 years). The nonlinear analysis was performed by the Phase Space Reconstruction method through a two-dimensional graph obtained by the Voice Analysis Program. The analysis was made by the 4-point crescent scale called curves-irregularity and spacing (C-IS). Statistical analysis employed the Mann-Whitney Test with a 5% significance level.

Results: PSR graphs showed open and closed trajectories across all lesions, with predominance of 4 curves (nodules ranged from 4 to 1 curves, while cysts and sulcus ranged from 4 to 2). Grade 1 (mild) irregularity predominated in nodules, cysts, and sulcus (42%, 43%, and 52%, respectively), with irregularities ranging from 0 to 3 degrees (normal to severe). Spacing was predominantly grade 1 (small) in nodules and cysts (46% and 48%, respectively), ranging from 0 to 3 degrees, while sulcus showed predominant grade 2 (medium) spacing (52%, ranging from 0 to 2 degrees). There was no significant difference ($p > 0.05$) between lesions on the C-IS scale.

Conclusion: Benign laryngeal lesions, including nodules, cysts, and sulcus, presented PSR graph with open and closed trajectories, predominance of 4 curves, mild irregularity and small to medium spacing.

Keywords. Voice. Dysphonia. Nonlinear Dynamics. Speech Acoustics. Speech, Language and Hearing Sciences

Acknowledgments. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001.

T4.2D - HOUSING DEMANDS FOR A SUCCESSFUL AGING-IN-PLACE: AN ANALYSIS THROUGH THE 5P'S MODEL

Janaina Moreno Garcia¹

janaina_mgarcia@usp.br

Carla da Silva Santana Castro¹

carla.santana@fmrp.usp.br

*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
Sãoocarlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

Summary

This study focuses on the concept of *aging-in-place* in terms of the functional, symbolic, and emotional bonds and meanings of homes and communities and how the environment is affecting well-being in old age." (WHO, 2017). **Objective:** To identify the demands of the elderly in relation to housing as a construct of AiP, based on the analysis of the 5 P's (Person, Place, Products, Services and Policies). **Methodology:** This is a retrospective, exploratory, cross-sectional, analytical and qualitative-quantitative study. 124 structured interview scripts were analyzed, about the demands of people over 50 years old from the main question "What current needs can influence your future?". The data were assessed through thematic content analysis. **Results:** Ninety-four participants were women, with a mean age of 67.7 years, married (52), with more than 10 years of schooling (61), 43 retired, and 48 participants continued to work. Two categories of analysis were listed: Housing demands, subdivided into Environment and Territory; and Demands of the elderly, which was subdivided into Personal and Third Factors. The demands of housing were expressed in the theme Built Environment, such as the need to adapt the residence; and in the Object Environment, such as the desire to purchase appliances. The demands of the elderly related to Personal Factors emerged concerns with health, finances and interpersonal relationships; and to the Third Parties he brought the knowledge of an active life in the community and in the family environment. **Conclusion:** Understanding these demands makes it possible to give voice and autonomy, favoring functional and active aging in the desired place.

Keywords: Aging in Place, Aging, Quality of Life and Autonomy

T4.3M - Health2Home - Mapping diagnostic imaging services at home and user needs: a scoping review

Giovana Fondato Costa¹

giovanafcosta@usp.br

Paula Costa Castro¹

castro@ufscar.br

*Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
Sãoocarlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil*

Abstract: Home diagnostic imaging services are increasingly popular due to their convenience and accessibility, particularly benefiting older adults, individuals with reduced mobility, and those living in remote areas. Customizing these services to individual needs is essential. However, the lack of guidelines compromises the quality and safety of care. This scoping review aims to map studies that identify the barriers and facilitators that influence the provision of home diagnostic imaging exams services. Several steps have been completed, including developing a search strategy using descriptors like diagnostic imaging, portable, home, and perceptions, and conducting searches in databases such as PubMed, Web of Science, Scopus, Embase, ACM Guide to Computing Literature, and Lilacs, without restrictions on publication date or language. A supplementary search was performed using Google Scholar. Two reviewers independently selected studies based on titles, abstracts, keywords, and full text, applying predefined inclusion criteria. The data analysis phase is currently underway. Quantitative data will be analyzed for mean, standard deviation, and frequency, while qualitative data will be categorized using content analysis. Partial results showed that of 1,471 studies identified, 25 were included, with 64% published after 2019 and covering 15 countries. The main diagnostic services studied include X-ray, electrocardiogram, electroencephalogram, and polysomnography. Expected outcomes include generating a list of design concept guidelines for home diagnostic imaging services aligned with user needs, along with identifying and categorizing barriers and facilitators of these services. These findings will support the Brazilian Health System (SUS) and companies in improving accessibility and implementation of these services.

Keywords. Diagnostic imaging; Portable; Residence; User-centered design.

Acknowledgments.

We acknowledge Interunit Graduate Program in Bioengineering, University of São Paulo and the Coordination for the Improvement of Higher Education Personnel - Brazil (CAPES), which financed this study (Financial Code 001).

T4.4D - Strategies for Minimizing the Effects of Observer Variability on Sagittal Parameter Measurements of the Spine

Adimilson dos Santos Delgado¹
delganeuro@usp.br

Bruna Souza Morais¹
b.souzamorais@usp.br

Helton Luiz Aparecido Defino²
hladefin@fmrp.usp.br

Arlindo Neto Montagnoli³
arlindo@ufscar.br

¹Interunit Graduate Program in Bioengineering, University of São Paulo, Av. Trabalhador
São-carlense, 400, Cx. P. 359, 13560-590, São Carlos - SP – Brazil

²Department of Biomechanics, Medicine and Rehabilitation of the Locomotor System, Faculty of
Medicine of Ribeirão Preto, University of São Paulo, Ribeirão Preto - SP – Brazil

³Department Electrical Engineering, Federal University of São Carlos, São Carlos - SP – Brazil

Abstract. Objective: Evaluate the influence of observer variability on thoracic kyphosis (TK) and lumbar lordosis (LL) measurements obtained with anatomical and functional spinal segmentation methods. **Background:** Spinal surgery planning typically relies on anatomical parameters. However, incorporating functional parameters that consider vertebrae orientation is important to minimizing surgical calculation errors. **Methods:** The authors developed parametric analysis software integrating traditional and functional methodologies. The proposed method included functional TK and functional LL calculated from the lines normal to the inflection points of the spine model. Using a synthetic lateral X-ray, the observer variability was computer-simulated by generating 20 landmark sets replicating 20 observers' annotations. The analysis also included 10 clinical X-rays, annotated twice by three judges with a one-week interval. Spinal curvature angles were derived using the anatomical and functional methods. Statistical analysis were performed. **Results:** For the synthetic X-ray, the proposed method presented significantly less variability: TK ($<\pm 2.5^\circ$, $p=0.00023$) and LL ($<\pm 5^\circ$, $p=0.00012$). For the clinical X-rays, interobserver reliability was higher for functional TK ($ICC>0.97$) and LL ($ICC>0.87$) compared to anatomical TK ($ICC<0.91$) and LL ($ICC<0.89$). Statistically significant differences were observed for both TK ($p=0.001$) and LL ($p=0.030$). Under the traditional method, observer variability led to measurement differences surpassing $\pm 19^\circ$, whereas differences with the proposed method were within $\pm 10^\circ$ for both parameters. **Conclusion:** The vertebral endplate is not the most suitable place to measure spinal sagittal curvatures. Small changes in landmark position significantly alter the measured Cobb angle. The proposed method offers a substantial advantage regarding the influence of observer variability.

Keywords. Sagittal alignment; Spinopelvic parameters; Vertebral parameters; Inflection point; Functional segmentation.